arXiv:1110.4878v1 [math.FA] 21 Oct 2011

L^2 dimensions of spaces of braid-invariant harmonic forms

Alexei Daletskii[a,*], Alexander Kalyuzhnyi[b]

[a]*Department of Mathematics, University of York, UK*
[b]*Institute of Mathematics, NAS, Kiev, Ukraine*

Abstract

Let X be a Riemannian manifold endowed with a co-compact isometric action of an infinite discrete group. We consider L^2 spaces of harmonic vector-valued forms on the product manifold X^N, which are invariant with respect to an action of the braid group B_N, and compute their von Neumann dimensions (*the braided* L^2*- Betti numbers*).

Keywords: Braid group, von Neumann algebra, L^2-Betti number, Configuration space
2010 MSC: Primary 58A10, 20F36, 46L10; Secondary 58A12, 58J22
Subject classification: Global Analysis, Analysis on Manifolds

---

1. Introduction

In recent years, there has been a growing interest in the analysis and geometry on various spaces of configurations (i.e. finite or infinite countable subsets) in a given topological space X, motivated by widespread applications in various parts of mathematics, including topology, representation theory, statistical mechanics, quantum field theory, mathematical biology, etc. (see [19], [33], [21], [20], [26], [9] and an extensive bibliography in [18], [8] and [31]). The development of the general mathematical framework for suitable classes of configurations is an important and challenging research theme, even in the case of finite number of components (particles) in each configuration. Many properties of the spaces of infinite configurations are in turn studied via the limit transition in the number of particles.

Observe that functions on the space of N-particle configurations in X can be identified with the functions on the Cartesian product X^N of N copies of X, which are invariant with

---

[*] Corresponding author.

Email addresses: `ad557@york.ac.uk` (Alexei Daletskii), `kalyuz@imath.kiev.ua` (Alexander Kalyuzhnyi)

respect to the natural action of the symmetric group S_N by permutations of arguments. Let us also note that certain spaces of mappings on X^N (e.g. vector-valued functions and differential forms in the case where X possesses a manifold structure) may admit more complicated groups of symmetries. An important example of an extended symmetry group is given by the braid group B_N, and the spaces of B_N-invariant differential forms on X^N can be regarded as spaces differential forms on "*braided configuration spaces*", which are related to the study of multi-particle systems obeying anyon and plekton statistics ([22], [16], [23]).

An important and intriguing question in this framework is the relationship between the properties associated with the multi-particle structure of these spaces, and the topology and geometry of the underlying space X. In many interesting cases, X is a non-compact Riemannian manifold of infinite volume, which makes it natural to discuss the above questions in the framework of the theory of L^2 invariants. This programme has been started in [15], [13], [14], [1].

Initiated by M. Atiyah, the theory of L^2 invariants of non-compact manifolds possessing infinite discrete groups of isometries serves as a brilliant example of application of von Neumann algebras in topology, geometry and mathematical physics. The initial framework introduced in [11] is as follows. Let X be a non-compact Riemannian manifold admitting an infinite discrete group G of isometries such that the quotient $K = X/G$ is compact. Then G acts by isometries on the spaces $L^2\Omega^m(X)$ of square-integrable m-forms over X. The projection $P^{(m)}$ onto the space $\mathcal{H}^m(X)$ of square-integrable harmonic m-forms (the m-th L^2-cohomology space) commutes with the action of G and thus belongs to the commutant $M_m(X)$ of this action, which is (under certain conditions) a II_∞-factor. The corresponding von Neumann trace $\mathrm{Tr}_{M_m} P^{(m)} =: \beta_m(X)$ gives a regularized dimension of the space $\mathcal{H}^m(X)$ and is called the L^2-Betti number of X (or K). Since the pioneering work [11], L^2-Betti numbers and other L^2 invariants have been studied by many authors (see e.g. [30] and references given there).

The construction above covers in an obvious way the case of the product manifold X^N endowed with the natural action of the product group G^N. The commutant $M_m(X^N)$ of the corresponding action in the space $L^2\Omega^m(X^N)$ contains the von Neumann subalgebra $\bigoplus_{k_1+\cdots+k_N=m} M_{k_1}(X) \otimes ... \otimes M_{k_N}(X)$, and the Künneth formula shows that

$$\beta_m(X^N) = \sum_{k_1+\cdots+k_N=m} \beta_{k_1}(X) \dots \beta_{k_N}(X).$$

The situation becomes much more complicated if we consider the spaces of forms on X^N admitting additional symmetries, for instance, invariant with respect to the action of the symmetric group S_N by permutations of arguments. In this case, the projection $\mathbf{P}_N^{(m)}$ onto the space of S_N-invariant harmonic forms *does not* in general belong to

$M_m(X^N)$.The minimal extension $\left\{M_m(X^N), \mathbf{P}_N^{(m)}\right\}''$ has been studied in [15], [13], [14], [1], where the corresponding trace formulae were derived and applied to the computation of L^2-Betti numbers of finite and infinite configuration spaces.

In the present paper, we consider the space $L^2_U\Omega^m(X^N, A_N)$ of square-integrable m-forms on X^N, which take values in a finite dimensional Abelian algebra A_N and are invariant with respect to a special representation U of the braid group B_N. The representation U is constructed as the tensor product of a given unitary representation π of B_N in A_N and the natural action $\mathfrak{S}$ of B_N in $L^2\Omega^m(X^N)$ by permutations.

Let $\mathbf{P}_N^{(m)}$ be the projection onto the space $\mathcal{H}^m_U(X^N, A_N)$ of harmonic forms in $L^2_U\Omega^m(X^N, A_N)$. The construction of the (minimal) von Neumann algebra containing $\mathbf{P}_N^{(m)}$ and computation of the trace of $\mathbf{P}_N^{(m)}$ involves the study of the von Neumann algebra

$$\left\{M_m(X^N) \otimes \mathrm{End}(A_N), U\right\}''. \tag{1}$$

We give a description of these von Neumann algebras and study the space $\mathcal{H}^m_U(X^N, A_N)$ in the situation where $A_N = (\mathbb{C}^2)^{\otimes N}$ and the representation π is generated by a 4×4 matrix C satisfying the braid equation

$$(C \otimes 1)(1 \otimes C)(C \otimes 1) = (1 \otimes C)(C \otimes 1)(1 \otimes C).$$

Representations of this type have been extensively studied, see [28].

The structure of the paper is as follows. In Sections 2 and 3, we derive general formulae for the traces on von Neumann algebras of the type (1). In Sections 4 and 5, we compute the L^2 dimensions of the spaces $\mathcal{H}^m_U(X^N, A_N)$. In Section 6, we define the space $\bigoplus_{N\in\mathbb{N}} \mathcal{H}^m_U(X^N, A_N)$ of "*harmonic m-forms on the braided configuration space*" and compute the *supertrace* of the corresponding projection (via the limit transition $N \to \infty$) in terms of the Euler characteristic of the underlying manifold X and certain invariant of the representation U.

In what follows we refer to [12], [32], [27] for general notions of the theory of von Neumann algebras. We denote by Tr_K the faithful normal semifinite trace on a II_∞- factor K.

2. Trace formula

We start with the review of some well-known facts from the theory of braid groups that we will need in what follows, see e.g. [10], [29]. The braid group B_N is generated by elements $b_1, \ldots, b_{N-1}$ satisfing the commutation relations

$$\begin{aligned} b_i b_j &= b_j b_i, \ |i-j| \geq 2, \\ b_i b_j b_i &= b_j b_i b_j, \ |i-j| = 1. \end{aligned}$$

The symmetric group S_N is a quotient group of B_N with respect to the relations $b_i^2 = 1$, $i = 1, \dots N-1$. We denote by $\phi : B_N \to S_N$ the corresponding canonical homomorphism.

Let us consider the *pure braid* group $B_N^{pure} := \operatorname{Ker} \phi$ (or the group of *color braids*). Define the forgetful homomorphism $f_{N-1} : B_N^{pure} \to B_{N-1}^{pure}$ which acts on a pure N-braid by forgetting its last string. Observe that the kernel of f_{N-1} is isomorphic to the free group F_{N-1}. Thus one has the split exact sequence of groups

$$1 \to F_{N-1} \to B_N^{pure} \to B_{N-1}^{pure} \to 1. \tag{2}$$

Thus the pure braid group can be decomposed into the semidirect product $B_N^{pure} = F_{N-1} \rtimes B_{N-1}^{pure}$ of the free group F_{N-1} and the group B_{N-1}^{pure} of pure $N-1$ braids. Iterating the decomposition procedure one can represent B_N^{pure} as the iterated semidirect product of free groups $F_1 \simeq \mathbb{Z}$, $F_2, \ \dots, F_{N-1}$ (the Artin normal form of B_N^{pure}). The generators $x_{i,j}$, $1 \leq i \leq j$, of the free group F_j, $j \leq N-1$, can be represented in the form

$$x_{i,j} = b_j b_{j-1} \dots b_{i+1} b_i^2 b_{i+1}^{-1} \dots b_{j-1}^{-1} b_j^{-1}.$$

Therefore the elements $x_{i,j}$, $1 \leq i \leq j \leq N-1$, generate B_N^{pure}.

In what follows, we will use the following special representation of the group B_N. Let $\mathfrak{S}$ be a unitary representation of the symmetric group S_N in a separable Hilbert space H and π be a unitary representation of B_N in a finite dimensional Hilbert space V. Then these representations generates a unitary representation U of the group B_N in Hilbert space $H \otimes V$ by the formula

$$U(b_i)(\xi \otimes v) = \mathfrak{S}(\phi(b_i))\xi \otimes \pi(b_i)v, \tag{3}$$

where $\xi \in H, \ v \in V$. A direct computation shows that

$$U(b_i)U(b_{i+1})U(b_i) = U(b_{i+1})U(b_i)U(b_{i+1}),$$

which implies that formula (3) indeed defines a unitary representation of the group B_N.

Let us introduce the subspace

$$(H \otimes V)_U = \{f \in H \otimes V | \, U(b_i)f = f, i = 1, \dots, N-1\} \tag{4}$$

of U-invariant elements of $H \otimes V$ and consider the corresponding orthogonal projection

$$P_U : H \otimes V \to (H \otimes V)_U \, .$$

Let M be a II_∞ factor acting in H and define $Q = M \otimes \operatorname{End}(V)$. In general, the projection P_U does not in general belong to Q. In what follows, we will extend Q to a suitable II_∞ factor containing P_U. We suppose that the following condition holds.

Condition 2.1. *The operators* $\mathfrak{S}(\sigma)$*,* $\sigma \in S_N$*,* $\sigma \neq e$*, do not belong to* M**.**

Condition 2.1 will be verified in the particular situations considered in the following sections. It implies that the formula

$$\alpha_\sigma(x) = \mathfrak{S}(\sigma) x \mathfrak{S}^*(\sigma), \quad x \in M \tag{5}$$

defines a nontrivial outer action

$$S_N \ni \sigma \mapsto \alpha_\sigma \in \mathrm{Aut}(M)$$

of the group S_N by automorphisms of the II_∞-factor M. Consider the corresponding cross-product $M \rtimes_\alpha S_N$. It is well known that an automorphism of a factor is free iff it is outer. Therefore (see e.g. [27], Proposition 1.4.4) $M' \cap (M \rtimes_\alpha S_N) = \mathbb{C}$ so that $M \rtimes_\alpha S_N$ is a (II_∞-) factor.

Lemma 2.2. *The map*

$$i : M \rtimes_\alpha S_N \ni (x, \sigma) \mapsto x U(\sigma) \in \{M, \{\mathfrak{S}(\sigma)\}_{\sigma \in S_N}\}'' \tag{6}$$

is an isomorphism of factors.

Proof. Observe that the map (6) is a non-trivial surjective normal homomorphism. It is well known that factors do not contain proper weakly closed two-sided ideals, which immediately implies that any homomorphisms between them is either identically zero or injective, and the result follows. ☐

Let us define the space

$$V^\pi = \{v \in V \mid \pi(\beta) v = v \text{ for all } \beta \in B_N^{pure}\}$$

of $\pi(B_N^{pure})$-invariant elements of V.

Lemma 2.3. *(i) The space* V^π *is invariant under the representation* π *of the braid group* B_N*.*

(ii) The restriction of the representation π *of* B_N *to the space* V^π *defines a representation* $\tilde{\pi}$ *of the symmetric group* S_N *via the formula*

$$\tilde{\pi}(\varphi(b_i)) = \pi(b_i) \restriction V^\pi, i = 1, \ldots, N-1.$$

Proof. (i). It is sufficient to show that the subspace V^π is invariant under the action of operators $\pi(b_i)$, $i = 1\dots, N-1$. Let us suppose that V^π is not invariant w.r.t. $\pi(b_{N-1})$. Then there exists an element $w \in \pi(b_{N-1})V^\pi$ orthogonal to V^π. Let $w = \pi(b_{N-1})v$, where $v \in V^\pi$. The equality $\pi(b^2_{N-1})v = v$ implies that $\pi(b_{N-1})w = v$ and $\pi(b^2_{N-1})w = w$. Then

$$v = \pi(b_{N-1})\pi(b^2_{N-2})\pi(b^{-1}_{N-1})v = \pi(b_{N-1})\pi(b^2_{N-2})w,$$

so that we have $\pi(b^2_{N-2})w = w$. The relations $b_{N-1}b_k = b_k b_{N-1}$, $k \le N-3$, and $\pi(x_{i,j})v = v$ imply that

$$\pi(x_{i,k})w = \pi(x_{i,k})\pi(b_{N-1})v = \pi(b_{N-1})\pi(x_{i,k})v = \pi(b_{N-1})v = w$$

for all $1 \le i \le k \le N-3$. We have also

$$\begin{aligned}\pi(x_{i,N-1})w &= \pi(b_{N-1})\pi(x_{i,N-2})\pi(b^{-1}_{N-1})w \\ &= \pi(b_{N-1})\pi(x_{i,N-2})v = \pi(b_{N-1})v = w\end{aligned}$$

because $\pi(b^{-1}_{N-1})w = v$ then for all $i = 1, \dots, N-2$. Finally, let us show that the vector w is invariant with respect to $\pi(x_{i,N-2})$, $i = 1, \dots, N-2$. The equality

$$\pi(b_{N-1})\pi(x_{i,N-2})\pi(b^{-1}_{N-1})v = v$$

implies that $\pi(b_{N-1})\pi(x_{i,N-2})w = v$, and

$$\pi(x_{i,N-2})w = \pi(b^{-1}_{N-1})v = w.$$

We see that w is invariant with respect to all operators $\pi(x_{i,j})$, $1 \le i \le j \le N-1$. Elements $x_{i,j}$ generate the pure braid group, which implies that $w \in V^\pi$**.** Therefore the space V^π is $\pi(b_{N-1})$-invariant under the operator
By similar arguments one can show that V^π is invariant w.r.t. the action of operators $\pi(b_k)$**,** $k = 1, \dots N-2$**.**

(ii) follows from the equality $\pi(b_i^2) \restriction V^\pi = \mathrm{id}_{V^\pi}$ and *(i)*. ☐

Consider the projection

$$p_\pi : V \to V^\pi.$$

We have the following result.

Theorem 2.4. *The projection $P_U : H \otimes V \mapsto (H \otimes V)_U$ can be represented in the form*

$$P_U = \frac{1 \otimes p_\pi}{N!} \sum_{\sigma \in S_N} \mathfrak{S}(\sigma) \otimes \tilde{\pi}(\sigma), \tag{7}$$

where $\tilde{\pi}$ is the unitary representation of the group S_N constructed in Lemma 2.3.

Proof. Let us denote by A the right-hand side of (7). **I**t follows from Lemma 2.3 that the relation $\pi(b_i)p_\pi = p_\pi\pi(b_i)$ holds for any $i = 1, \dots, N-1$. Therefore $A \in \mathrm{End}\,(H \otimes V)$ is a projection.

It is sufficient to show that images of P and A coincide. It is obvious that if $\xi \in (H \otimes V)_U$ then $(1 \otimes p_\pi)\xi = \xi$ and $\mathfrak{S}_\sigma \otimes \tilde{\pi}(\sigma)\xi = \xi$. Therefore the space $(H \otimes V)_U = P(H \otimes V)$ is contained in the image of A. Let us prove the converse inclusion. Let an element $\xi \in H \otimes V$ belong to the image of A. Then $\xi \in H \otimes V^\pi$ and $\mathfrak{S}(\sigma) \otimes \tilde{\pi}(\sigma)\xi = \xi$. Since $\pi(b_i) \restriction V^\pi = \tilde{\pi}(\varphi(b_i))$ we have

$$U(b_i)\xi = \mathfrak{S}(\varphi(\mathfrak{b}_\mathfrak{i})) \otimes \pi(b_i)\xi = \mathfrak{S}_{\varphi(b_i)} \otimes \tilde{\pi}(\varphi(b_i))\xi = \xi,$$

i. e. ξ is U-invariant. Thus the image of A is contained in the space $(H \otimes V)_U = P_U(H \otimes V)$. ☐

Corollary 2.5. *We have the inclusion* $(H \otimes V)_U \subset H \otimes V^\pi$**.**

Let us introduce the von Neumann algebra $Q_U := \{Q, \{U(\beta)\}_{\beta \in B_N}\}''$ generated by Q and the operators $U(\beta), \beta \in B_N$. It is obvious that $P_U \in Q_U$.

Lemma 2.6. *Von Neumann algebras* $(M \rtimes S_N) \otimes \mathrm{End}(V)$ *and* Q_U *are isomorphic.*

Proof. Recall that operators $\mathfrak{S}(\phi(\beta))$, $\beta \in B_N$, $\beta \neq e$, do not belong to M, which implies that operators

$$U(\beta) = \mathfrak{S}(\phi(\beta)) \otimes \pi(\beta),\ \beta \in B_N,\ \beta \notin B_N^{pure},$$

do not belong to Q. Therefore the representation U generates a nontrivial outer action α' of the group S_N on the II_∞ factor Q. Applying the arguments similar to the proof of Lemma 2.2 one can see that there exists an isomorphism

$$(Q \rtimes_{\alpha'} S_N) \simeq \{Q, \{U(\beta)\}_{\beta \in B_N}\}''. \tag{8}$$

On the other hand, for any $\beta \in B_N$ we have the inclusion $\pi(\beta) \in \mathrm{End}(V)$**,** which implies that the factors $Q \rtimes_{\alpha'} S_N$ and $(M \rtimes_\alpha S_N) \otimes \mathrm{End}(V)$ are isomorphic, and the result follows. ☐

Corollary 2.7. Q_U *is a* II_∞*-factor.*

Proof. Follows from the fact that$(M \rtimes S_N) \otimes \mathrm{End}(V)$ is a II_∞-factor. ☐

Corollary 2.8. *It follows from (2.6) that* $\mathrm{Tr}_{Q_U} = \mathrm{Tr}_{M \rtimes S_N} \times \mathrm{tr}_{\mathrm{End}(V)}$.

Without loss of generality we assume that the trace on $\mathrm{End}(V)$ is normalized, that is, $\mathrm{tr}_{\mathrm{End}(V)}(1) = 1$.

Corollary 2.9. *For any $A \in M$ and $B \in \mathrm{End}\,(V)$ commuting with p_π we have*

$$\mathrm{Tr}_{Q_U}((A \otimes B)P_U) = \frac{\mathrm{Tr}_M(A)}{N!}\mathrm{tr}_{\mathrm{End}(V)}(Bp_\pi). \tag{9}$$

Proof. It follows the properties of the trace on a semidirect product of factors that for any $\sigma \in S_N$ the following equality holds:

$$\begin{aligned} \mathrm{Tr}_{Q_U}(A\mathfrak{S}(\sigma) \otimes Bp_\pi\tilde{\pi}(\sigma)) &= \mathrm{Tr}_{M \rtimes S_N}(A\mathfrak{S}(\sigma))\mathrm{tr}_{\mathrm{End}(V)}(Bp_\pi\tilde{\pi}(\sigma)) \\ &= \delta_{e,\sigma}\mathrm{Tr}_M(A)\mathrm{tr}_{\mathrm{End}(V)}(Bp_\pi\tilde{\pi}(\sigma)), \end{aligned}$$

where $\delta_{e,\sigma}$ is the Kronecker symbol. Formula (9) follows now from (7). ☐

Remark 2.1. Introduce the projection

$$P_s = \frac{1}{N!}\sum_{\sigma \in S_N} \mathfrak{S}_\sigma.$$

It follows from (9) that the following formula holds:

$$\begin{aligned} \mathrm{Tr}_{Q_U}((A \otimes B)P_U) &= \mathrm{Tr}_M(P_s)\mathrm{Tr}_{Q_U}((A \otimes B)(1 \otimes p_\pi)) \\ &= \frac{1}{N!}\mathrm{Tr}_{Q_U}((A \otimes B)(1 \otimes p_\pi)). \end{aligned}$$

In order to extend formula (9) to general elements of Q_U, we introduce II_∞-factors

$$Q^1 := M \otimes \mathrm{End}\,(V^\pi) \text{ and } Q^2 := M \otimes \mathrm{End}\,((V^\pi)^\perp).$$

It is well known that any finite dimensional unitary representation of a locally compact group is completely reducible. Thus the representation π can be decomposed into a direct sum $\pi = \pi^1 \oplus \pi^2$ where $\pi^1 = \tilde{\pi} = \pi \restriction V^\pi$ and $\pi^2 = \pi \restriction (V^\pi)^\perp$. Observe that in the case when π is irreducible we have $V^\pi = \{0\}$ and π^1 is a zero representation. Therefore we can introduce von Neumann algebras

$$\tilde{Q}^1 = \{Q^1, \{\mathfrak{S}(\sigma) \otimes \tilde{\pi}(\sigma)\}_{\sigma \in S_N}\}'' \text{ and } \tilde{Q}^2 = \{Q^2, \{\mathfrak{S}(\phi(\beta)) \otimes \pi^2(\beta)\}_{\beta \in B_N}\}''.$$

Clearly $\tilde{Q}^1$ and $\tilde{Q}^2$ are isomorphic to II_∞-factors $Q^1 \rtimes S_N$ and $Q^2 \rtimes S_N$ respectively, and $P_U \in \tilde{Q}^1$.

We define the following natural trace on the von Neumann algebra $\tilde{Q}^1 \oplus \tilde{Q}^2$ setting

$$\mathrm{Tr}_{\tilde{Q}^1 \oplus \tilde{Q}^2}(A_1 \oplus A_2) = \frac{\dim V^\pi}{\dim V}\mathrm{Tr}_{\tilde{Q}^1}(A_1) + \frac{\dim\,(V^\pi)^\perp}{\dim V}\mathrm{Tr}_{\tilde{Q}^2}(A_2), \tag{10}$$

$A_1 \in \tilde{Q}^1$, $A_2 \in \tilde{Q}^2$.

Consider an arbitrary $A \in Q$ commuting with $1 \otimes p_\pi$. Then $A \in Q^1 \oplus Q^2$ and $AP_U \in \tilde{Q}^1$. Observe that $\frac{\dim V^\pi}{\dim V} = \mathrm{tr}_{\mathrm{End}(V)}(p_\pi)$ and $A(1 \otimes p_\pi) \in Q^1$. Applying the arguments similar to the proof of Corollary 2.9 one can see that

$$\mathrm{Tr}_{\tilde{Q}^1 \oplus \tilde{Q}^2}(AP_U) = \frac{\mathrm{Tr}_{Q^1}(A(1 \otimes p_\pi))}{N!} \mathrm{tr}_{\mathrm{End}(V)}(p_\pi). \tag{11}$$

Remark 2.2. Note that formula (11) is compatible with (9). Indeed, if $A \otimes B \in M \otimes \mathrm{End}\ (V)$ and $B \in \mathrm{End}\ (V)$ commutes with p_π, then $A \otimes B \in Q^1 \oplus Q^2$ and by (11) we have

$$\begin{aligned} \mathrm{Tr}_{\tilde{Q}^1 \oplus \tilde{Q}^2}((A \otimes B)P_U) &= \frac{\mathrm{Tr}_{Q^1}(A \otimes Bp_\pi)}{N!} \mathrm{tr}_{\mathrm{End}(V)}(p_\pi) \\ &= \frac{\mathrm{Tr}_M(A)}{N!} \mathrm{tr}_{\mathrm{End}(V^\pi)}(Bp_\pi) \mathrm{tr}_{\mathrm{End}(V)}(p_\pi) \\ &= \frac{\mathrm{Tr}_M(A)}{N!} \mathrm{tr}_{\mathrm{End}(V)}(Bp_\pi) \end{aligned}$$

because the traces $\mathrm{tr}_{\mathrm{End}(V)}$ and $\mathrm{tr}_{\mathrm{End}(V^\pi)}$ are normalized.

3. Braided multi-particle spaces and the corresponding trace

In this section we apply the general construction discussed above to the case where the space H has the form of the tensor product $H_0^{\otimes N}$ of N copies of a separable Hilbert space H_0 and

$$S_N \ni \sigma \mapsto \mathfrak{S}(\sigma) \in \mathrm{End}(H_0^{\otimes N}) \tag{12}$$

is the natural action of S_N in $H_0^{\otimes N}$ by permutations, so that $\mathfrak{S}(\phi(b_i))$ is the permutation of the i-th and $i+1$-th components in $H_0^{\otimes N}$. As in the previous section, the representation U is defined by formula (3), that is,

$$U(b_i)(\xi \otimes v) = \mathfrak{S}(\phi(b_i))\xi \otimes \pi(b_i)v, \tag{13}$$

where $\xi \in H,\ v \in V$, elements $b_1, \ldots, b_{N-1}$ are the generators of the group B_N and $\phi : B_N \to S_N$ is the canonical homomorphism. The corresponding subspace $(H \otimes V)_U$ of invariant elements (cf. (4)) will be called the *braided N-particle space.*

Let M_0 be a II_∞ factor acting in H_0 and define the corresponding II_∞ factors $M = M_0^{\otimes N}$ and $Q = M \otimes \mathrm{End}\, V$ acting in $H = H_0^{\otimes N}$ and $H \otimes V$.

The following result was proved in [13] (see also [14]).

Proposition 3.1. *Representation $\mathfrak{S}$ defined by formula (12) satisfies Condition 2.1, that is,* $\mathfrak{S}(\sigma) \notin M$ *for all* $\sigma \in S_N$.

Proposition 3.1 implies that we can apply the theory developed in the previous section. According to Lemma 2.6 we have

$$P_U \in Q_U = (M_0^{\otimes N} \times_\alpha S_N) \otimes \mathrm{End}(V). \tag{14}$$

In what follows we will show that Q_U is the minimal extension of $M_0^{\otimes N}$ containing and P_U.

Theorem 3.2. *Factors* Q_U *and* $Q_P := \{Q, P_U\}''$ *are isomorphic.*

Proof. The inclusion $Q_P \subset (M_0^{\otimes N} \times_\alpha S_N) \otimes \mathrm{End}(V)$ follows from (14).

Let us prove the inverse inclusion. Observe that operators $\pi(b_i) \in \mathrm{End}(V)$ and hence $1_{M_0^{\otimes N} \times_\alpha S_N} \otimes \pi(b_i) \in Q_P$. This it suffices to prove that the operators $\mathfrak{S}(\phi(b_i)) \otimes 1_{\mathrm{End}(V)}$ belong to Q_P. Denote $P_i = \frac{1}{2}(1 + \mathfrak{S}(\phi(b_i)) \otimes 1_{\mathrm{End}(V)})$. The II_∞-factor M can be represented in a form $N \otimes B(l_2)$, where N is a II_1 factor and $B(l_2)$ is a space of all bounded operators in the Hilbert space $l_2 = l_2(\mathbb{N})$. Therefore M contains the isometry $v = 1_N \otimes T$, where $Te_i = e_{i+1}$ is the operator of unilateral shift in l_2. It is clear that $(v^*)^m \to 0$, $m \to \infty$, strongly. Therefore we have strong convergence

$$\begin{aligned} &\left((1 \otimes 1 \otimes (v^*)^m \otimes \cdots \otimes (v^*)^m) \otimes 1_{\mathrm{End}(V)}\right) \cdot \\ &\cdot P\left((1 \otimes 1 \otimes v^m \otimes \cdots \otimes v^m) \otimes 1_{\mathrm{End}(V)}\right) \to \frac{2}{N!} P_1, \end{aligned} \tag{15}$$

$m \to \infty$. Thus $P_1 \in M$, which implies that $P_1 \in M \otimes \mathrm{End}(V) \subset Q_P$. Similar arguments show that $P_i \in Q_P$ for any $i = 1, \ldots, N-1$. □

Remark 3.1. It follows from Theorem 3.2 that formulae (9) and (11) are valid for Tr_{Q_P} instead of Tr_{Q_U}.

4. Von Neumann dimensions of the spaces of braid-invariant harmonic forms

Let X be a smooth connected Riemannian manifold admitting an infinite discrete group G of isometries such that the quotient $K = X/G$ is a compact connected Riemannian manifold.

In this section we consider the De Rahm complex of square-integrable forms on X^N taking values in a finite dimensional Abelian algebra A_N [24]. We apply the trace formula (9) in order to find the von Neumann dimensions of the spaces of U-invariant harmonic forms (the *braided* L^2*- Betti numbers*).

4.1. *Setting: Von Neumann algebras associated with infinite coverings of compact manifolds*

Let us describe the framework introduced by M. Atiyah in his theory of L^2-Betti numbers, which we will use during the rest of the paper. For a detailed exposition, see [11] and e.g. [30].

We assume that there exists an infinite discrete group G acting freely on X by isometries and that $K = X/G$ is a compact Riemannian manifold. That is,

$$G \to X \to K \tag{16}$$

is a Galois (normal) cover of K.

Throughout this section, we fix $m = 1, ..., d,\ d = \dim X - 1$, and use the following general notations:
$L^2\Omega^m(X)$ - the space of square-integrable m-forms on X;
$L^2\Omega^m(K)$ - the space of square-integrable m-forms on K;
$H_X^{(m)}$ - Hodge-de Rahm Laplacian on X (considered as a self-adjoint operator in $L^2\Omega^m(X)$);
$\mathcal{H}^m(X) := \operatorname{Ker} H_X^{(m)}$ - the space of square-integrable harmonic m-forms on X.

The action of G in X generates in the natural way the unitary representation of G in $L^2\Omega^m(X)$, which we denote

$$G \ni g \mapsto T_g \in \operatorname{End}(L^2\Omega^m(X)). \tag{17}$$

Let M_m be the commutant of this representation,

$$M_m = \{T_g\}'_{g\in G} \subset \operatorname{End}(L^2\Omega^m(X)). \tag{18}$$

It is clear that the space $L^2\Omega^m(X)$ can be described in the following way:

$$L^2\Omega^m(X) = L^2\Omega^m(K) \otimes l^2(G), \tag{19}$$

with the group action obtaining the form

$$T_g = id \otimes L_g, \tag{20}$$

$g \in G$, where $L_g,\ g \in G$, are operators of the left regular representation of G. Then

$$\operatorname{End}(L^2\Omega^m(X)) = \operatorname{End}(L^2\Omega^m(K)) \otimes \operatorname{End}(l^2(G)) \tag{21}$$

and

$$M_m = \operatorname{End}(L^2\Omega^m(K)) \otimes \mathcal{R}(G), \tag{22}$$

where $\mathcal{R}(G)$ is the von Neumann algebra generated by the right regular representation of G.

In what follows, we assume that G is an ICC group, that is,

$$\text{\textit{all non-trivial classes of conjugate elements are infinite.}} \tag{23}$$

This ensures that $\mathcal{R}(G)$ is a II_1-factor (see e.g. [32]). Thus M is a II_∞-factor.

Let us consider the orthogonal projection

$$P^{(m)}_{Har} : L^2\Omega^m(X) \to \mathcal{H}^m(X) \tag{24}$$

and its integral kernel

$$k(x, y) \in T_xX \otimes T_yX. \tag{25}$$

Then, because of the G-invariance of the Laplacian $H^{(m)}_X$, we have $P_{Har} \in M_m$. It was shown in [11] that

$$\mathrm{Tr}_M P^{(m)}_{Har} = \int_K \mathrm{tr}\, k(x, x)\, dx, \tag{26}$$

where tr is the usual matrix trace and dx is the Riemannian volume on K. Let us remark that, because of G-invariance, $k(m, m)$ is a well-defined function on K. Moreover, it is known that $H^{(m)}_X$ is elliptic regular, which implies that the kernel k is smooth. Thus

$$\beta_m := \mathrm{Tr}_M\ P^{(m)}_{Har} < \infty. \tag{27}$$

The numbers $\beta_m,\ m = 0, 1, ..., d$, are called the L^2-Betti numbers of X (or K) associated with G. L^2-Betti numbers are homotopy invariants of K ([17]). It is known that

$$\sum_{m=0}^{d} (-1)^m \beta_m = \chi(K), \tag{28}$$

where $\chi(K)$ is the Euler characteristic of K (L^2 index theorem [11]).

Observe that all of the above can be applied to the product group G^N acting on the product manifolds X^N (instead of G acting on X). The Künneth formula

$$\mathcal{H}^m(X^N) = \bigoplus_{m_1+\cdots+m_N=m} \mathcal{H}^{m_1}(X) \otimes \cdots \otimes \mathcal{H}^{m_N}(X). \tag{29}$$

implies that the corresponding L^2-Betti numbers of X^N have the form

$$\beta_m(X^N) = \sum_{m_1+\cdots+m_N=m} \beta_{m_1} \dots \beta_{m_N}. \tag{30}$$

4.2. L^2-dimensions of the spaces of braid-invariant harmonic forms

Let A_N be a finite dimensional Abelian algebra possessing a structure of a finite dimensional Hilbert space. Consider the Hilbert space $L^2\Omega^m(X^N) \otimes A_N$ of square integrable (with respect to the volume measure) A_N-valued m-forms on X^N.

Recall that the group G acts on X by isometries. Denote by

$$M_m(X^N) \subset \mathrm{End}(L^2\Omega^m(X^N))$$

and

$$Q^{(m,N)} := M_m(X^N) \otimes \mathrm{End}(A_N) \subset \mathrm{End}(L^2\Omega^m(X^N) \otimes A_N)$$

the commutant of the induced natural action of G^N (by isometries) on $L^2\Omega^m(X^N)$ and $L^2\Omega^m(X^N) \otimes A_N$, respectively. Here $\mathrm{End}(A_N)$ is the space of endomorphisms of A_N with respect to its Hilbert space structure (but not in general morphisms of the algebra A_N). Both $M_m(X^N)$ and $Q^{(m,N)}$ are II_∞ factors.

We adopt the construction of Section 2 with $H = L^2\Omega^m(X^N)$ and $V = A_N$. Consider a unitary representation

$$\pi : B_N \ni b \mapsto \pi(b) \in \mathrm{End}(A_N)$$

of the braid group B_N in A_N, and the corresponding representation (cf. (3))

$$U : B_N \ni b \mapsto U(b) = \mathfrak{S}(\phi(b)) \otimes \pi(b) \in \mathrm{End}(L^2\Omega^m(X^N) \otimes A_N)$$

in the space $L^2\Omega^m(X^N) \otimes A_N$, where $\mathfrak{S}$ is the natural action the group S_N in $L^2\Omega^q(X^N)$ defined in local coordinates by the formula

$$\mathfrak{S}(\sigma)f((x_m)_{m=1,\dots N})dx_{i_1} \wedge \dots \wedge dx_{i_q} = f((x_{\sigma(m)})_{m=1,\dots N})dx_{\sigma(i_1)} \wedge \dots \wedge dx_{\sigma(i_m)}. \quad (31)$$

Here $\sigma \in S_N$ is the permutation $(1, ..., N) \mapsto (\sigma(1), ..., \sigma(N))$. Let

$$L^2\Omega^m_U(X^N, A_N) := \left(L^2\Omega^m(X^N) \otimes A_N\right)_U$$

be the Hilbert space of U-invariant elements of $L^2\Omega^m(X^N) \otimes A_N$ (cf. (4)), and consider the corresponding projection

$$P_U^{(m)} : L^2\Omega^m(X^N) \otimes A_N \to L^2\Omega^m_U(X^N, A_N).$$

Let $A_N^\pi := \{a \in A_N : \pi(b)a = a,\ b \in B_N^{pure}\}$ be the space of all $\pi(B_N^{pure})$-invariant elements of A_N. In what follows we will assume that

$$A_N^\pi \text{ is a subalgebra of } A_N. \quad (32)$$

The latter condition holds in all the examples considered in Section 5 below.

Let $\Omega^m(X^N)$ be the space of smooth differential m-forms on X^N, and

$$d^m_{X^N} : L^2\Omega^m(X^N) \to L^2\Omega^{m+1}(X^N),\ \mathrm{Dom}(d^m_{X^N}) = \Omega^m(X^N)$$

be the Hodge differential on X^N. It can be extended to an operator

$$\mathbf{d}^m_N := d^m_{X^N} \otimes 1_{A_N} : L^2\Omega^m(X^N) \otimes A_N \to L^2\Omega^{m+1}(X^N) \otimes A_N,$$

where 1_{A_N} is the identity operator on A_N.

Proposition 4.1.

(i) The exterior product of two U-invariant A_N-valued forms is also U-invariant.

(ii) The differential $\mathbf{d}^m_N$ preserves the space of U-invariant forms, that is,

$$\mathbf{d}^m_N \left(P^{(m)}_U \left(\Omega^m(X^N) \otimes A_N \right) \right) \subset P^{(m+1)}_U \left(\Omega^{m+1}(X^N) \otimes A_N \right).$$

Proof. (i) The statement follows from (7) and condition (32)**.**

(ii) The statement follows from the fact that $\mathbf{d}^m_{X^N}$ commutes with operators $\mathfrak{S}(\phi(b_i)) \otimes id$ and $id \otimes \pi(b_i)$, $i = 1, \ldots, N-1$. □

Let us consider the Hodge-de Rham Laplacian $H^{(m)}_{X^N}$ and define

$$\mathbf{H}^{(m,N)} := H^{(m)}_{X^N} \otimes 1_{A_N}.$$

$\mathbf{H}^{(m,N)}$ is a selfajoint operator in $L^2\Omega^m(X^N) \otimes A_N$. It follows from the definition of the operators $\mathbf{d}^m_N$ and $\mathbf{H}^{(m,N)}$ that

$$\mathbf{H}^{(m,N)} = \mathbf{d}^m_N \left(\mathbf{d}^m_N\right)^* + \left(\mathbf{d}^{m-1}_N\right) \mathbf{d}^{m-1}_N.$$

Proposition 4.1 implies that $\mathbf{H}^{(m,N)}$ preserves the space of U-invariant forms. Moreover, $\mathbf{H}^{(m,N)}$ commutes with the operators $U(b)$, $b \in B_N$.

Let us consider the projection

$$P^{(m,N)}_{Har} : L^2\Omega^m(X^N) \to \mathcal{H}^m(X^N).$$

Introduce the space

$$\mathcal{H}^m_U(X^N, A_N) := \left(\mathcal{H}^m(X^N) \otimes A_N\right)_U,$$

of U-invariant elements of $\mathcal{H}^m(X^N) \otimes A_N$. Observe that the projection

$$P_{Har}^{(m,N)} \otimes 1_{A_N} : L^2\Omega^m(X^N) \otimes A_N \to \mathcal{H}^m(X^N) \otimes A_N$$

commutes with the projection

$$P_U^{(m,N)} : L^2\Omega^m(X^N) \otimes A_N \to L^2\Omega_\pi^m(X^N, A_N).$$

Therefore the operator

$$\mathbf{P}^{(m,N)} := (P_{Har}^{(m,N)} \otimes 1_{A_N})P_U^{(m,N)} : L^2\Omega^m(X^N) \otimes A_N \to \mathcal{H}_U^m(X^N, A_N).$$

is the projection, too.

Our goal is to compute the L^2-dimension of the space $\mathcal{H}_U^m(X^N, A_N)$, that is, the von Neumann trace of the projection $\mathbf{P}^{(m,N)}$. Observe that $P_{Har}^{(m,N)} \in M_m(X^N)$, which implies that $P_{Har}^{(m,N)} \otimes 1_{A_N} \in Q^{(m,N)}$.

Proposition 4.2. *The factor $M_m(X^N)$ does not contain the operators $\mathfrak{S}(\phi(b))$, $b \in B_N$, $b \notin B_N^{pure}$.*

Proof. Note that the space $L^2\Omega^m(X^N)$ can be represented in the form

$$L^2\Omega^m(X^N) = L^2(X)^{\otimes N} \otimes \Lambda^m(\mathbb{C}^N),$$

where $\Lambda^m(\mathbb{C}^N)$ is a m-th direct summand of the exterior algebra $\Lambda(\mathbb{C}^N)$. Then the factor $M_m(X^N)$ has the form $M_m(X^N) = M^{\otimes N} \otimes \operatorname{End}(\Lambda^m(\mathbb{C}^N))$, where M is the commutant of the action of the group G in $L^2(X)$. The representation $\mathfrak{S}$ of the group S_N can be decomposed into the product $\mathfrak{S} = \mathcal{S} \otimes \mathcal{S}'$, where $\mathcal{S}$ and $\mathcal{S}'$ are natural actions of S_N by permutations in $L^2(X)^{\otimes N}$ and $\Lambda^m(\mathbb{C}^N)$, respectively (cf. (12) and (31)). It follows from Proposition 3.1that $\mathcal{S}(\sigma) \notin M^{\otimes N}$, $\sigma \neq e$. Therefore $\mathfrak{S}(\sigma) \notin M_m(X^N)$, $\sigma \neq e$, which implies the result. ☐

Thus Condition 2.1 is satisfied, and we can apply the results of Section 2. Let

$$Q_P^{(m,N)} := \{Q^{(m,N)}, P_U^{(m,N)}\}''$$

be the minimal von Neumann algebra containing $Q^{(m,N)}$ and $P_U^{(m,N)}$.

Theorem 4.3. *The von Neumann algebras $Q_P^{(m,N)}$, $\big(M_m(X^N) \rtimes S_N\big) \otimes \operatorname{End}(A_N)$ and $\{Q^{(m,N)}, \{U(b)\}_{b \in B_N}\}''$ are isomorphic.*

Proof. The first isomorphism of can be proved by the arguments similar to the proof of Theorem 3.2. The second isomorphism follows from Lemma 2.6. ☐

Theorem 4.3 limplies that the von Neumann algebra $Q_P^{(m,N)}$ is a II_∞ factor (since $\big(M_m(X^N) \rtimes S_N\big) \otimes \mathrm{End}\ (A_N)$ is so).

Theorem 4.4. *The projection $\mathbf{P}^{(m,N)}$ belongs to $Q_P^{(m,N)}$ and its trace is given by the formula*

$$\mathrm{Tr}_{Q_P^{(m,N)}}(\mathbf{P}^{(m,N)}) = \frac{\mathrm{Tr}_{\mathrm{End}(A_N)}(p_\pi)}{N!} \sum_{m_1+\cdots+m_N=m} \beta_{m_1} \ldots \beta_{m_N} \tag{33}$$

Here β_m, $m = 0, 1, \ldots, \dim X$, are the L^2-Betti numbers of X and p_π is the projection onto the subalgebra A_N^π of B_N^{pure}-invariant elements of A_N.

Proof. The inclusion $P^{(m,N)} \in Q_P^{(m,N)}$ follows from the definition of the von Neumann algebra $Q_P^{(m,N)}$. The equality (33) follows from Corollary 2.9 with $A \otimes B = P_{Har}^{(m,N)} \otimes 1_{A_N}$ and the Künneth formula

$$\mathcal{H}^m(X^N) = \bigoplus_{m_1+\cdots+m_N=m} \mathcal{H}^{m_1}(X) \otimes \cdots \otimes \mathcal{H}^{m_N}(X). \tag{34}$$

☐

We will use the notation $\mathbf{b}_m(X^N) = \mathrm{Tr}_{Q_P^{(m,N)}}(\mathbf{P}^{(m,N)})$ and call $\mathbf{b}_m(X^N)$ the m-th *braided L^2-Betti number of X^N.*

Let us introduce the constant

$$C_N^\pi = \frac{\mathrm{tr}_{\mathrm{End}(A_N)}(p_\pi)}{N!}.$$

Observe that the trace on the algebra $\mathrm{End}(A_N)$ is normalized, that is, $\mathrm{tr}_{\mathrm{End}(A_N)}(1_{A_N}) = 1$, which implies that

$$C_N^\pi = \frac{\dim A_N^\pi}{N! \dim A_N} \leq \frac{1}{N!}. \tag{35}$$

Then formula (33) can be rewritten in the form

$$\mathbf{b}_m(X^N) = C_N^\pi \ \mathrm{Tr}_{M^{(m)}}(P_{Har}^{(m,N)}). \tag{36}$$

Remark 4.1. In the situation where the representation π reduces to a representation of the symmetric group S_N (i.e. $\pi(b_i^2) = 1$) the projection $p_\pi = 1$ and formula (33) can be rewritten in a form

$$\mathrm{Tr}_{Q_\pi^m}(P_{Har}^{(m,N)}) = \frac{1}{N!}\mathrm{Tr}_{M^{(m)}}(P_{Har}^{(m,N)})$$

In this case the braided L^2-Betti numbers b_m do not depend on the algebra A_N and coincide with the L^2-Betti numbers of the space of N-point configurations $\Gamma_N(X)$ of X ([13], [14]).

5. Examples

In this section we consider examples associated with particular representations of the braid group B_N. We set $A_N = (\mathbb{C}^2)^{\otimes N}$ ($= \mathbb{C}^{2^N}$) equipped with the natural Abelian algebra structure.

An important class of representations π can be constructed in the following way. Let C be a complex unitary 4×4-matrix satisfying the braid equation:

$$(C \otimes 1)(1 \otimes C)(C \otimes 1) = (1 \otimes C)(C \otimes 1)(1 \otimes C), \tag{37}$$

where 1 denotes the identity 2×2-matrix. For all generators $b_i \in B_N$ define an operator $\pi(b_i)$ acting in A_N by the formula

$$\pi(b_i) := 1 \otimes 1 \otimes \cdots \otimes C \otimes \cdots \otimes 1, \tag{38}$$

where C acts on the i-th and $i+1$-th components of the tensor product $(\mathbb{C}^2)^{\otimes \mathbb{N}}$. Then π defines a unitary representation of B_N. This representation is not reduced to a representation of S_N if and only if $C^2 \neq 1$.

Remark 5.1. Representations (38) are extensively studied. It is known that the solutions of the equation (37) are in one-to-one correspondence with constant solutions R of the Yang-Baxter equation

$$R^{k_1k_2}_{j_1j_2} R^{l_1k_3}_{k_1j_3} R^{l_2l_3}_{k_2k_3} = R^{k_2k_3}_{j_2j_3} R^{k_1l_3}_{j_1k_3} R^{l_1l_2}_{k_1k_2}. \tag{39}$$

More precisely, $C = R\,\Sigma$, where Σ is a numerical matrix with elements $\Sigma^{ij}_{kl} = \delta^j_k \delta^i_l$ and δ^j_k denotes the Kröneker symbol. All two-dimensional solutions of (39) have been classified in [25], which gives the complete list of unitary 4×4-matrices satisfying the braid equation (37).

Example 1. Let C be an arbitrary unitary solution of the equation (37) and $C^2 = 1$. For instance,

$$C = \begin{pmatrix} 0 & 0 & 0 & q \\ 0 & \varepsilon & 0 & 0 \\ 0 & 0 & \varepsilon & 0 \\ \overline{q} & 0 & 0 & 0 \end{pmatrix},$$

where $\varepsilon = \pm 1$ and $|q| = 1$. Then the representation π is reduced to a representation of S_N , so that the corresponding representation $\tilde{\pi}$ of the pure braid group B_N^{pure} is trivial ($\tilde{\pi}(b) = id$ for any $b \in B_N^{pure}$) and thus $A_N^{\pi} = A_N$. Then $C_N^{\pi} = \frac{1}{N!}$ and

$$\mathbf{b}_m(X^N) = \frac{1}{N!} \sum_{m_1 + \cdots + m_N = m} \beta_{m_1} \ldots \beta_{m_N}. \tag{40}$$

Example 2. Let $q \in \mathbb{C}$ be such that $|q| = 1$ and $q^2 \neq 1$. Set

$$C = \begin{pmatrix} 0 & 0 & 0 & q \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ q & 0 & 0 & 0 \end{pmatrix},$$

Then C is unitary solution of (37) and $C^2 \neq 1$.

Proposition 5.1. $\dim A_N^\pi = 2$ *for any* $N \in \mathbb{N}$.

Proof. A direct calculation using formula (38). ☐

We can apply the results of Section 4 and compute the braided L^2-Betti numbers $\mathbf{b}_m(X^N)$ of a connected cocompact Riemannian manifold X. Formula (35) implies that $C_N^\pi = \frac{1}{2^{N-1}N!}$. Therefore

$$\mathbf{b}_m(X^N) = \frac{1}{2^{N-1}N!} \sum_{m_1+\cdots+m_N=m} \beta_{m_1} \ldots \beta_{m_N}, \tag{41}$$

where β_i, $i = 0, 1, 2$, are the L^2-Betti numbers of X.

It this example, we can give an explicit description of the braided N-particle space. A direct calculation shows that the only non-zero components of $f = (f_1, \ldots, f_{2^N}) \in (H \otimes V)_U$ are $f_{a_{N-1}}$ and f_{a_N}, where the sequence a_N is defined recursively as follows: $a_1 = 2$ and $a_N = 2^N - a_{N-1} + 1$. Moreover $f_{a_{N-1}}$ and f_{a_N} are symmetric elements, so that $(H \otimes V)_U = H_0^{\hat{\otimes} N} \otimes \mathbb{C}^2$, where $\hat{\otimes}$ denotes the symmetric tensor product.

Example 3. Let $q \in \mathbb{C}$ be such that $|q| = 1$ and $q^2 \neq 1$. Set

$$C = \begin{pmatrix} q & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

Then C is unitary solution of (37) and $C^2 \neq 1$.

Proposition 5.2. $dim\ A_N^\pi = N + 1$.

Proof. Let $e_0 = (1, 0)^t$, $e_1 = (0, 1)^t$ be the canonical basis in $\mathbb{C}^2$. Denote by $e_{i_1 \ldots i_n} = e_{i_1} \otimes \cdots \otimes e_{i_N}$ the corresponding basis in $\mathbb{C}^{2^N}$.

Consider first the case of $N = 2$. We have $B_2 = \{b\}$ and $\pi(b)e_{00} = qe_{00}$, $\pi(b)e_{01} = e_{10}$, $\pi(b)e_{10} = e_{01}$, $\pi(b)e_{11} = e_{11}$. Therefore the basis of A^{π}_N consists of e_{01}, e_{10} and e_{11}, and $\dim A^{\pi}_N = 3$.

A direct check shows that $e_{\underbrace{11\ldots11}_{N}}$ and $e_{\underbrace{11\ldots10}_{N}}$ belong to A^{π}_N. Let us prove that for any multiindex $I = i_1 \ldots i_N$, $i_k \in \{0,1\}$, such that $e_I \in A^{\pi}_N$ the equality $i_N = 0$ implies that $i_1 = \ldots i_{N-1} = 1$. Indeed, let $i_N = 0$ and $k := \max_{j\in\{1,2,\ldots,N-1\}}\{i_j = 0\} > 0$. Then it is easy to see that

$$\begin{aligned} e_I = \pi(x_{k,N-1})e_I &= \pi(b_{N-1}b_{N-2}\ldots b_{k+1}b_k^2 b_{k+1}^{-1}\ldots b_{N-2}^{-1}b_{N-1}^{-1})e_I \\ &= \pi(b_{N-1}b_{N-2}\ldots b_{k+1}b_k^2)e_{i_1\ldots i_{k-1}001\ldots1} = q^2 e_I, \end{aligned}$$

because the space A^{π}_N is invariant under the representation π and $\pi(b_k^2)e_{i_1\ldots i_{k-1}001\ldots1} = q^2 e_{i_1\ldots i_{k-1}001\ldots1}$ (recall that $\pi(b_i) = 1\otimes\cdots\otimes1\otimes C\otimes1\otimes\cdots\otimes1$). Therefore $e_I = 0$.

Let us prove that $e_I \in A^{\pi}_N$ iff $e_{I1} \in A^{\pi}_{N+1}$. Indeed, let $e_I \in A^{\pi}_N$. To prove that $e_{I1} \in A^{\pi}_{N+1}$ it suffice to check that $\pi(x_{k,N})e_{I1} = e_{I1}$ for all $k \le N$. If $i_N = 1$ we have $\pi(x_{k,N})e_{I1} = \pi(b_N x_{N-1,k})e_{I1} = \pi(b_N)e_{I1} = e_{I1}$ since $\pi(x_{k,N-1})e_I = e_I$. If $i_N = 0$ then by previous statement we have that $i_1 = \cdots = i_{N-1} = 1$ and $\pi(x_{k,N})e_{I1} = \pi(b_N x_{k,N-1})e_{11\ldots10} = \pi(b_N)e_{11\ldots10} = e_{11\ldots101} = e_{I1}$.

The equalities $\pi(x_{i,j})e_{I1} = e_{I1}$, $i \le j \le N-1$ and the fact that the operators $\pi(x_{i,j})$ are acting only in first $N-1$ indexes imply that $e_I \in A^{\pi}_N$ provided $e_{I1} \in A^{\pi}_{N+1}$. This statement combined with the induction arguments and the fact that $e_{\underbrace{11\ldots10}_{N+1}} \in A^{\pi}_{N+1}$ complete the proof. ☐

It follows from Proposition 5.2 that $C^{\pi}_N = \frac{N+1}{2^N N!}$. Therefore the braided L^2-Betti numbers $\mathbf{b}_m(X^N)$ of X have the following form:

$$\mathbf{b}_m(X^N) = \frac{N+1}{2^N N!}\sum_{m_1+\cdots+m_N=m}\beta_{m_1}\ldots\beta_{m_N}, \tag{42}$$

where β_i, $i = 0,1,2$, are the L^2-Betti numbers of X.

The structure of the corresponding braided N-particle space is more complicated then is previous example. It is not hard to see that the representation $\tilde{\pi}$ of the symmetric group in the space A^{π}_N that corresponds to the representation π can be decomposed into a direct sum of two cyclic representations $\tilde{\pi}_1$ and $\tilde{\pi}_2$. Representation $\tilde{\pi}_1$ is trivial and acts in one-dimensional space generated by the basis element $e_{11\ldots1}$ while representation $\tilde{\pi}_2$ permutetes the other N basis elements in A^{π}_N. Therefore the space $H^{\otimes_\pi N}$ can be decomposed into the direct sum of the space $L^2(X)^{\hat{\otimes} N}$ of symmetric functions and the space $H^{\otimes_{\tilde{\pi}_2} N}$ of elements from $L^2(X^N) \otimes \mathbb{C}^N$ that are invariant under representation $\tilde{U}_N$ of the group S_N given by the relation

$$(\tilde{U}_N(\sigma_i)f)(x_1, \dots, x_i, x_{i+1}, \dots x_N) = \tilde{\pi}_2(\sigma_i)f(x_1, \dots, x_{i+1}, x_i, \dots, x_N),$$

where $\sigma_1, \dots, \sigma_{N-1}$ are generating elements of the group S_N.

Example 4. Let $q \in \mathbb{C}$ be such that $|q| = 1$ and $q^2 \neq 1$

$$C = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & q & 0 & 0 \\ 0 & 0 & q & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}.$$

Then C is unitary solution of (37) and $C^2 \neq 1$.

Proposition 5.3. $\dim A_N^\pi = 0$.

Proof. Similar to the Example 2 we can show that $\pi(b_k^2)e_{i_1 \dots i_k i_{k+1} \dots i_N} = q^2 e_{i_1 \dots i_N}$ iff $i_k i_{k+1} = 01$ or $i_k i_{k+1} = 10$. Therefore any vector $v \in A_N^\pi$ should have the form $v = \alpha e_{00\dots0} + \beta e_{11\dots1}$. On the other hand, $\pi(x_{1,2})e_{00\dots0} = q^2 e_{00\dots0}$ and $\pi(x_{1,2})e_{11\dots1} = q^2 e_{11\dots1}$, so that $\pi(x_{1,2})v = q^2 v$, which implies that $v = 0$. Therefore $\dim A_N^\pi = 0$. ☐

Thus, $\mathbf{b}_k(X^N) = 0$.

6. Fock space of braid-invariant harmonic forms: L^2-dimensions and index

Let us consider the infinite disjoint union

$$\mathfrak{X} = \bigsqcup_{N=1}^{\infty} X^N.$$

The space $L^2\Omega^m(\mathfrak{X})$ of square-integrable m-forms on $\mathfrak{X}$ has the form

$$L^2\Omega^m(\mathfrak{X}) = \bigoplus_N L^2\Omega^m(X^N). \tag{43}$$

We can define the Hodge-de Rham Laplacian $H_{\mathfrak{X}}^{(m)}$ on $L^2\Omega^m(\mathfrak{X})$ component-wise, that is,

$$H_{\mathfrak{X}}^{(m)} := \sum_N H_{X^N}^{(m)}.$$

For the corresponding spaces of harmonic forms we obviously have the decomposition

$$\mathcal{H}^m(\mathfrak{X}) := \mathrm{Ker}\ H_{\mathfrak{X}}^{(m)} = \bigoplus_N \mathcal{H}^m(X^N). \tag{44}$$

Formulae (43) and (44) motivate us to *define* the space

$$L^2\Omega_U^m(\mathfrak{X}{,}A) := \bigoplus_N L^2\Omega_U^m(X^N, A_N),$$

which can be regarded as the space of ”*m-forms on the braided configuration space*”. Similar to the case of the de Rham complex (43), the Hodge-de Rham Laplacian $\mathbf{H}^{(m)}$ on $L_\pi^2\Omega^m(\mathfrak{X})$ can be defined component-wise,

$$\mathbf{H}^{(m)} := \sum_N \mathbf{H}^{(m,N)}.$$

The corresponding space of U-invariant harmonic m-forms can be decomposed into the direct sum

$$\mathcal{H}_U^m(\mathfrak{X}{,}A) := \mathrm{Ker}\ \mathbf{H}^{(m)} = \bigoplus_N \mathcal{H}_U^m(X^N, A_N).$$

Let $\mathbf{P} : = \sum_m \mathbf{P}^{(m)}$, where $\mathbf{P}^{(m)}$ is the projection onto $\mathcal{H}_U^m(\mathfrak{X}{,}A)$. Moreover, we have the equality $\mathbf{P}^{(m)} = \sum_N \mathbf{P}^{(m,N)}$, which implies that $\mathbf{P}^{(m)}$ an element of the von Neumann algebra $Q_P^{(m)} := \bigoplus\limits_N Q_P^{(m,N)}$.

We introduce a regularized dimension $\mathbf{b}_m(\mathfrak{X})$ of the space $\mathcal{H}_U^m(\mathfrak{X}{,}A)$ by the formula

$$\mathbf{b}_m(\mathfrak{X}) = \sum_{N=0}^{\infty} \mathbf{b}_m^{(N)} = \mathrm{Tr}_{Q_P^{(m)}}\ \mathbf{P}^{(m)}$$

and define the *supertrace*

$$\mathbb{STR}\ \mathbf{P} = \sum_{m=0}^{\infty} (-1)^m \mathbf{b}_m(\mathfrak{X}). \tag{45}$$

Here we use the convention $\mathbf{b}_0 = 1$.

Theorem 6.1. *The series in the right-hand side of (45) converges absolutely and*

$$\mathbb{STR}\ \mathbf{P} = \sum_N C_N^\pi \chi(K)^N < \infty,$$

where $\chi(K)$ is the Euler characteristic of K.

Proof. Formula (35) shows that $C_N^\pi \leq \frac{1}{N!}$. Then

$$\begin{aligned}
\sum_{m=0}^{\infty} \mathbf{b}_m(\mathfrak{X}) &\leq \sum_{m=0}^{\infty} \sum_{N=0}^{\infty} \frac{1}{N!} \sum_{k_1+k_2+\ldots+k_N=m} \beta_{k_1} \ldots \beta_{k_N} \\
&= \sum_{N=0}^{\infty} \frac{1}{N!} \left(\sum_k \beta_k \right)^N < \infty.
\end{aligned}$$

Then

$$\begin{aligned}
\mathbb{STR}\, \mathbf{P} &= \sum_{m=0}^{\infty} (-1)^n \, \mathbf{b}_m(\mathfrak{X}) \\
&= \sum_{m=0}^{\infty} (-1)^m \sum_N C_N^\pi \sum_{k_1+k_2+\ldots+k_N=m} \beta_{k_1} \ldots \beta_{k_N} \\
&= \sum_N C_N^\pi \sum_{k_1,k_2,\ldots,k_N} (-1)^{k_1} \beta_{k_1} \ldots (-1)^{k_N} \beta_{k_N} \\
&= \sum_N C_N^\pi \left(\sum_k (-1)^k \beta_k \right)^N \\
&= \sum_N C_N^\pi \chi(K)^N
\end{aligned}$$

because of the equality $\sum_m (-1)^m \beta_m = \chi(K)$ (the L^2 index theorem [11]). □

Let us consider the examples of the previous section. In all these examples we have $A_N = (\mathbb{C}^2)^{\otimes N}$ so that $\dim A_N = 2^N$. Thus, we can apply Theorem 6.1 and compute $\mathbb{STR}\, \mathbf{P}_\pi$.

Example 1 (revisited). $C_N^\pi = \frac{1}{N!}$ and $\mathbb{STR}\, \mathbf{P} = e^{-\chi(K)}$. The latter expression coincides with the formula derived in [1] for the case the de Rham complex over the configuration space equipped with the Poisson measure.

Example 2 (revisited). $C_N^\pi = \frac{1}{2^{N-1} N!}$. Then

$$\mathbb{STR}\, \mathbf{P} = \sum_N (-1)^N C_N^\pi \chi(K)^N = \sum_N (-1)^N \frac{2}{N!} \left(\frac{\chi(K)}{2} \right)^N = 2e^{-\frac{\chi(K)}{2}}.$$

Example 3 (revisited). $C_N^\pi = \frac{N+1}{2^N N!}$. Then

$$\begin{aligned}
\mathbb{STR}\,\mathbf{P} &= \sum_N (-1)^N C_N^\pi \chi(K)^N \\
&= \sum_N (-1)^N \frac{N}{N!}\left(\frac{\chi(K)}{2}\right)^N + \sum_N (-1)^N \frac{1}{N!}\left(\frac{\chi(K)}{2}\right)^N \\
&= -\frac{\chi(K)}{2}\sum_N (-1)^{N-1}\frac{1}{(N-1)!}\left(\frac{\chi(K)}{2}\right)^{N-1} + \sum_N (-1)^N \frac{1}{N!}\left(\frac{\chi(K)}{2}\right)^N \\
&= 1 - \frac{\chi(K)}{2} e^{-\frac{\chi(K)}{2}} + e^{-\frac{\chi(K)}{2}}.
\end{aligned}$$

Remark 6.1. Under the additional assumption of $\dim X = 2$, the only non-zero L^2-Betti number of X is $\beta = \beta_1(X)$. Then, according to formula (33), the only non-zero braided L^2-Betti number of X^N is $\mathbf{b}_N(X^N) = C_N^\pi \beta^N$, and therefore

$$\mathbf{b}_m(\mathfrak{X}) = C_N^\pi \beta^m,\ m = 1, 2, 3, \ldots.$$

Remark 6.2. The right-hand side of formula (45) can be understood as a regularized index of the Dirac operator $D + D^*$, where $D := \sum_m \sum_N \mathbf{d}_N^m$, see [11] and e.g. [30] for the discussion of von Neumann supertraces in geometry and topology of Riemannian manifolds and their relation to L^2-index theorems, and [1], [2], [3] for the extension of these notions to the framework of infinite configuration spaces.

Acknowledgments

The authors would like to thank Sergio Albeverio, Yuri Kondratiev, Eugene Lytvynov and Leonid Vainerman for helpful discussions.

References

References

[1] S. Albeverio, A. Daletskii, L^2-Betti numbers of Poisson configuration spaces, *Publ. RIMS* ***42*** (2006), 649-682.

[2] S. Albeverio, A. Daletskii, A. Kalyuzhnyj, Traces of semigroups associated with interacting particle systems, *J. Funct. Anal.* **246** (2007), 196-216.

[3] S. Albeverio, A. Daletskii, A. Kalyuzhnyj, Random Witten Laplasians: Traces of semigroups, L^2-Betti numbers and index, *J. Eur. Math. Soc. (JEMS)* **10** (2008), 571–599.

[4] S. Albeverio, A. Daletskii, Yu. Kondratiev, E. Lytvynov, Laplace operators in de Rham complexes associated with measures on configuration spaces, *J. Geom. Phys.* **47** (2003), 259-302.

[5] S. Albeverio, A. Daletskii, E. Lytvynov, Laplace operators on differential forms over configuration spaces, *J. Geom. Phys.* **37** (2001), 15–46.

[6] S. Albeverio, A. Daletskii, E. Lytvynov, De Rham cohomology of configuration spaces with Poisson measure, *J. Funct. Anal.* **185** (2001), 240-273.

[7] S. Albeverio, Yu. Kondratiev, M. Röckner, Analysis and geometry on configuration spaces, J. Funct. Anal. 154 (1998) 444–500.

[8] S. Albeverio, Yu. Kondratiev, M. Röckner, Analysis and geometry on configuration spaces: The Gibbsian case, J. Funct. Anal. 157 (1998) 242–291.

[9] S. Albeverio, Yu. Kondratiev, M. Röckner, Diffeomorphism groups and current algebras: Configuration spaces analysis in quantum theory, Rev. Math. Phys. 11 (1999) 1–23.

[10] E. Artin, Theorie der Zöpfe, Abh. Math. Sem. Hamburg Univ., 1926, **4**, p. 47–72.

[11] M. F. Atiyah, Elliptic operators, discrete groups and von Neumann algebras, *Colloque ”Analyse et Topologie” en l’Honneur de Henri Cartan* (Orsay, 1974), pp. 43–72. *Asterisque*, No. 32-33, Soc. Math. France, Paris, 1976.

[12] O.Bratelli, D. W. Robinson, Operator algebras and quantum statistical Mechanics, Springer-Verlag, 1979.

[13] A. Daletskii, A. Kalyuzhnyj, Permutations in tensor products of factors and L^2- Betti numbers of configuration spaces. Proceedings of V International Conference ”Symmetry in Nonlinear Mathematical Physics” (Kiev, June 2003), eds. A.G. Nikitin, V.M. Boyko, R.O. Popovych, I.A.Yehorchenko, *Proceedings of Institute of Mathematics of NAS of Ukraine* **50** (2004), 1025-1528.

[14] A. Daletskii, A. Kalyuzhnyj, Permutations in tensor products of factors and L^2 cohomology of configuration spaces, *Methods of Functional Analysis and Topology.*, **12** (2006), 341–352.

[15] A. Daletskii, Yu. Samilenko, Von Neumann dimensions of symmetric and antisymmetric tensor products, *Methods of Functional Analysis and Topology* **9** (2003), No. 2, 123-133.

[16] G. Dell'Antonio, R. Figari, A. Teta. Statistics in Space Dimension Two. — Lett. Math. Phys. **40**, 1997, 235–256.

[17] J. Dodziuk, De Rham-Hodge theory for L^p-cohomology of infinite coverings, *Topology* **16** (1977), 157–165.

[18] D.J. Daley, D. Vere-Jones, An Introduction to the Theory of Point Processes, vol. I: Elementary Theory and Methods, 2nd ed., Probab. Appl. (N.Y.), Springer, New York, 2003.

[19] E. R. Fadell, S. Y. Husseini, "Geometry and Topology of Configuration Spaces", Springer Verlag, Berlin/Neidelberg, 2001.

[20] H.O. Georgii, Canonical Gibbs Measures, Lecture Notes Math., vol. 760, Springer, Berlin, 1979.

[21] G.A. Goldin, J. Grodnik, R.T. Powers, D.H. Sharp, Nonrelativistic current algebra in the N/V limit, J. Math. Phys. 15 (1974) 88–100.

[22] G.A. Goldin, D. H. Sharp, Diffeomorphism groups, anyon fields, and q-commutators, Phys. Rew. Lett., **76** (1996) 1183–1187.

[23] G.A.Goldin, Sh. Majid, On the Fock space for nonrelativistic anyon fields and Braided tensor products, *J. Math. Phys.* **45** (2004).

[24] A. Guichardet, Cohomologie des groupes topologiques et des algebres de Lie, Cedic/Fernan Nathan, 1980.

[25] J. Hietarinta, Solving the two-dimensional constant quantum Yang-Baxter equation, *J. Math. Phys*. **34** (1993) 1725 – 1756.

[26] R.S. Ismagilov, Representations of Infinite-Dimensional Groups, Transl. Math. Monogr., vol. 152, Amer. Math. Soc., Providence, RI, 1996.

[27] V. Jones, S. Sunder, Introduction to subfactors, London Math. Soc. Lecture Note Series **234**, Cambridge Univ. Press.

[28] Ch. Kessel, Quantum groups, Graduate Texts in Math. **155**, Springer, 1995.

[29] Ch. Kassel, V. Turaev, Braid groups, Graduate Texts in Math. **247**, Springer, 2008.

[30] V. Mathai, L^2 invariants of covering spaces, *in* “Geometric analysis and Lie theory in mathematics and physics,” Austral. Math. Soc. Lect. Ser. **11**, pp. 209–242, Cambridge Univ. Press, Cambridge, 1998.

[31] M. Röckner, Stochastic analysis on configuration spaces: Basic ideas and recent results, in: J. Jost, et al. (Eds.), New Directions in Dirichlet Forms, Stud. Adv. Math., vol. 8, Amer. Math. Soc., Providence, RI, 1998, pp. 157–232.

[32] M. Takesaki, Theory of operator algebras, Springer-Verlag, 1979.

[33] A.M. Vershik, I.M. Gel’fand, M.I. Graev, Representations of the group of diffeomorphisms, Russian Math. Surveys 30 (6) (1975) 1–50.